

# Constraining the Dimensions of Interstellar Cold Clouds Encountered by the Heliosphere

Anonymous Author(s)

## ABSTRACT

The spatial dimensions of interstellar cold clouds that the Sun may encounter remain unknown, creating uncertainties spanning four orders of magnitude in heliosphere crossing durations (100 yr to 1 Myr). We develop a Bayesian framework combining ISM cloud size distributions, crossing time constraints, and cosmogenic  $^{10}\text{Be}$  detectability requirements to constrain cloud dimensions. Our posterior analysis yields a median cloud size of 2.09 pc with 68% credible interval [0.11, 10.0] pc. Monte Carlo propagation through the full model predicts a median crossing time of  $\sim 29,000$  years (16–84%: [2,900, 296,000] yr). For a proposed 2–3 Mya cold cloud encounter, the  $^{10}\text{Be}$  signal has an 89.3% detection probability in marine sediment records, with median SNR of 8.8 after radioactive decay correction. Heliosphere compression modeling shows that a cold cloud with density  $100 \text{ cm}^{-3}$  would shrink the heliosphere to  $\sim 12$  AU, enhancing cosmic ray flux by a factor of  $\sim 3.5$ . These results provide quantitative constraints for interpreting geological isotope records and planning future investigations.

## 1 INTRODUCTION

The heliosphere shields Earth from galactic cosmic rays (GCRs), but encounters with dense interstellar clouds can dramatically compress the heliosphere, exposing the inner solar system to enhanced radiation [4, 5]. Nica et al. [3] modeled cosmogenic  $^{10}\text{Be}$  production during such encounters but noted that cloud dimensions are unknown, necessitating a broad range of crossing times from 100 years to 1 Myr.

Constraining cloud dimensions is essential for: (1) predicting realistic crossing durations, (2) assessing  $^{10}\text{Be}$  signal detectability in geological archives, and (3) evaluating the biological and climatic impacts of enhanced cosmic ray exposure. The interstellar medium (ISM) near the Sun contains both warm and cold phases [1], with cold clouds ( $T \sim 20 \text{ K}$ ,  $n_H \sim 100 \text{ cm}^{-3}$ ) following size distributions governed by turbulent fragmentation [2].

## 2 METHODS

### 2.1 Cloud Size Distribution

We model cloud sizes using a log-normal distribution motivated by ISM turbulence, with mean  $\log_{10}(L/\text{pc}) = 0$  and standard deviation 1.0. The size-density relation follows Larson's scaling:  $n_H \propto L^{-0.7}$ .

### 2.2 Crossing Time Model

Crossing time depends on cloud size, solar velocity ( $v_\odot = 26 \text{ km/s}$ ), and impact parameter  $b$ :

$$t_{\text{cross}} = \frac{2R\sqrt{1-b^2}}{v_\odot} \quad (1)$$

where  $R$  is the cloud radius and  $b \in [0, 1]$  is uniformly distributed.

### 2.3 Bayesian Inference

The posterior on cloud size incorporates: (1) a log-normal ISM prior, (2) crossing time constraints (100 yr to 1 Myr), (3)  $^{10}\text{Be}$  detectability likelihood in marine sediments, and (4) ISM consistency.

### 2.4 $^{10}\text{Be}$ Signal Model

During cloud encounters, GCR flux is enhanced by  $\sim 4\times$ . For a 2.5 Mya event, signal decay reduces detection by a factor  $\exp(-\ln 2 \cdot 2.5 \times 10^6 / 1.39 \times 10^6) \approx 0.29$ .

## 3 RESULTS

### 3.1 Bayesian Cloud Dimensions

The posterior median cloud size is 2.09 pc (68% CI: [0.11, 10.0] pc), corresponding to a median crossing time of  $\sim 29,000$  years (Table 1).

Table 1: Cloud dimension and crossing time estimates.

Parameter	Median	68% CI
Cloud size (pc)	1.03	[0.11, 10.0]
Crossing time (yr)	29,000	[2,900, 296,000]
$^{10}\text{Be}$ SNR (marine)	8.8	—
Detection probability	89.3%	—




Figure 1: (a) Prior, likelihood, and posterior distributions for cloud size. (b) Implied crossing time distribution.

### 3.2 Heliosphere Response

A cold cloud with  $n_H = 100 \text{ cm}^{-3}$  compresses the heliosphere from 120 AU to  $\sim 12$  AU, enhancing GCR flux by  $\sim 3.5\times$  (Figure 2).

### 3.3 $^{10}\text{Be}$ Detectability

$^{10}\text{Be}$  signals become detectable (SNR > 2) for crossing times exceeding  $\sim 500$  years in marine sediments (Figure 3). Ice cores provide better sensitivity but are limited to more recent events.




Figure 2: (a) Heliosphere radius versus ISM density. (b) GCR flux enhancement.




Figure 3:  $^{10}\text{Be}$  signal-to-noise ratio versus crossing duration for marine sediments and ice cores.

### 3.4 Known Cloud Analysis

The Local Interstellar Cloud ( $\sim 3$  pc) would produce a  $\sim 113,000$ -year crossing. The Local Leo Cold Cloud ( $\sim 5$  pc, 23 pc away) could be encountered in  $\sim 0.86$  Myr.

## 4 CONCLUSION

Our Bayesian framework constrains interstellar cold cloud dimensions to a median of  $\sim 1\text{--}2$  pc with crossing times of  $\sim 10^4\text{--}10^5$  years. The high detection probability (89%) in marine sediments for a 2–3 Mya event supports the feasibility of identifying past cloud encounters through  $^{10}\text{Be}$  anomalies. These constraints narrow the uncertainty from four to approximately two orders of magnitude, providing actionable predictions for targeted searches in geological archives.

## 5 LIMITATIONS AND ETHICAL CONSIDERATIONS

Cloud size estimates depend on the assumed ISM turbulence model and may not apply to all cloud environments. The spherical cloud approximation underestimates crossing time variability for filamentary structures. Solar velocity uncertainties propagate directly into crossing time estimates. This fundamental astrophysics research has no direct ethical implications.

## REFERENCES

- [1] Priscilla C. Frisch, Seth Redfield, and Jonathan D. Slavin. 2011. The Interstellar Medium Surrounding the Sun. *Annual Review of Astronomy and Astrophysics* 49 (2011), 237–279.
- [2] Richard B. Larson. 1981. Turbulence and Star Formation in Molecular Clouds. *Monthly Notices of the Royal Astronomical Society* 194 (1981), 809–826.
- [3] Sami Nica et al. 2026. Modeling Cosmogenic  $^{10}\text{Be}$  During the Heliosphere's Encounter with an Interstellar Cold Cloud. *arXiv preprint arXiv:2601.07983* (2026).
- [4] Klaus Scherer et al. 2006. Interstellar-Terrestrial Relations: Variable Cosmic Environments, the Dynamic Heliosphere, and Their Imprints on Terrestrial Archives and Climate. *Space Science Reviews* 127 (2006), 327–465.
- [5] Gary P. Zank et al. 2013. Heliospheric Structure: The Bow Wave and Beyond. *The Astrophysical Journal* 763 (2013), 20.

117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232